

Clinical Indicator Diseases for Adult HIV infection

	AIDS-defining conditions	Other conditions where HIV testing should be offered
Respiratory	Tuberculosis Pneumocystis	Bacterial pneumonia Aspergillosis
Neurology	Cerebral toxoplasmosis Primary cerebral lymphoma Cryptococcal meningitis Progressive multifocal Leucoencephalopathy	Aseptic meningitis /encephalitis Cerebral abscess Space occupying lesion of unknown cause Guillain–Barré syndrome Transverse myelitis Peripheral neuropathy Dementia Leucoencephalopathy
Dermatology	Kaposi’s sarcoma	Severe or recalcitrant seborrhoeic dermatitis Severe or recalcitrant psoriasis Multidermatomal or recurrent herpes zoster
Gastroenterology	Persistent Cryptosporidiosis	Oral candidiasis, Oral hairy leukoplakia Chronic diarrhoea of unknown cause Weight loss of unknown cause Salmonella, shigella or campylobacter Hepatitis B and C infections
Oncology	Non-Hodgkin’s lymphoma	Anal cancer or anal intraepithelial dysplasia Lung cancer Seminoma Head and neck cancer Hodgkin’s lymphoma Castleman’s disease
Gynaecology	Cervical cancer	Vaginal intraepithelial neoplasia Cervical intraepithelial neoplasia Grade 2 or above
Haematology		Any unexplained blood dyscrasia including: thrombocytopenia, neutropenia lymphopenia
Ophthalmology	Cytomegalovirus retinitis	Infective retinal diseases including herpesviruses and toxoplasma Any unexplained retinopathy
ENT		Lymphadenopathy of unknown cause Chronic parotitis Lymphoepithelial parotid cysts
Other		Mononucleosis-like syndrome (primary HIV infection) Pyrexia of unknown origin Any lymphadenopathy of unknown cause Any sexually transmitted Infection